


CHARGER™ 0.035" Balloon Dilatation Catheter

COUNT ON IT.



Charger™ is the 0.035" balloon you can count on.

Boston Scientific's Charger is a steadfast 0.035" Balloon Dilatation Catheter designed to deliver results in both routine and challenging cases. It offers excellent push and track, a low profile, rapid deflation, non-compliant dilatation, and a full range of sizes.



Post-Stent Dilatation

- Non-compliant for excellent stent-to-wall apposition
- Higher pressure when needed (up to 20 atm for 5 × 200 mm)



Low Profile


- 0.040" tip over a 0.035" wire
- 6 × 150 and 7 × 80 through 5 Fr
- 10 × 80 through 6 Fr
- Excellent rewrap



Broad Size Matrix

- Balloon lengths 20–200 mm
- Diameters 3–12 mm
- Shafts 75, 135 cm

Charger™ 0.035" Balloon Dilatation Catheter



CHARGER™ OTW

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Directions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions.

INTENDED USE/INDICATIONS FOR USE: The Charger Balloon Dilatation Catheter is indicated for Percutaneous Transluminal Angioplasty (PTA) in the peripheral vasculature, including iliac, femoral, popliteal, tibial, peroneal, subclavian, and renal arteries and for the treatment of obstructive lesions of native or synthetic arteriovenous dialysis fistulae. The Charger Balloon Dilatation Catheter is also indicated for post-dilatation of balloon expandable and self-expanding stents in the peripheral vasculature. **CONTRAINDICATIONS:** None Known. **WARNINGS:** The Charger PTA balloon catheter is not for use in the coronary arteries. **PRECAUTIONS:** The Charger PTA Balloon Dilatation Catheter shall only be used by physicians trained in the performance of percutaneous transluminal angioplasty. The Charger PTA Balloon Dilatation Catheter should be used with caution for procedures involving calcified lesions or synthetic vascular grafts due to the abrasive nature of these inflation sites. The Charger PTA Balloon Dilatation Catheters are not intended for injection of contrast medium. Precautions to prevent or reduce clotting should be taken when any catheter is used: • Consider systemic anticoagulation. • Flush or rinse all products with sterile saline or a similar solution prior to use. **ADVERSE EVENTS:** The complications that may result from a balloon dilatation procedure include, but are not limited to: • Allergic reaction (device, contrast medium and medications) • Arteriovenous fistula • Embolization (air, device, plaque, etc.) • Hematoma • Hemorrhage, including bleeding at puncture site • Pseudoaneurysm • Sepsis/infection • Thromboembolic episodes • Vessel injury, e.g. dissection, perforation, rupture • Vessel occlusion • Vessel spasm **REV AA**

Charger is a registered trademark of Boston Scientific Corporation or its affiliates. All other trademarks are property of their respective owners.

Boston Scientific
Advancing science for life™

Peripheral Interventions

300 Boston Scientific Way
Marlborough, MA 01752-1234
www.bostonscientific.com

To order product or for more information contact customer service at 1.888.272.1001.

© 2015 Boston Scientific Corporation or its affiliates. All rights reserved.

PI-28511-AB MAR2015